arXiv:hep-ph/0204168v1 15 Apr 2002

Yaroslavl State University
Preprint YARU-HE-02/02

hep-ph/0204168

Electron mass operator in a strong magnetic field

A.V. Kuznetsov, N.V. Mikheev and M.V. Osipov

Division of Theoretical Physics, Department of Physics,
Yaroslavl State University, Sovietskaya 14,
150000 Yaroslavl, Russian Federation

Abstract

The electron mass operator in a strong magnetic field is calculated by summation of the leading log contributions in all orders of the perturbation theory. An influence of the strong field on the virtual photon polarization operator is taken into account. The contribution of higher Landau levels of virtual electrons, along with the ground Landau level, is shown to be essential in the leading log approximation.

Keywords: electron mass operator; strong magnetic field asymptotics; rainbow Feynman diagram

PACS numbers: 11.10.Jj, 12.20.Ds, 14.60.Cd

Submitted to Modern Physics Letters A

Investigations of the asymptotic properties of diagrams and operators of the quantum electrodynamics in strong magnetic fields $B \gg B_e$ ($B_e = m_e^2/e \simeq 4.41 \cdot 10^{13}$ G [1]) is of conceptual interest both from the standpoint of the searches of the borders of the perturbation theory applicability, and in view of possible astrophysical applications. The researches of this type are being performed by many authors during a rather long time. For example, a history of calculations of the electron mass operator in a strong magnetic field lasts more than 30 years already. However, as we show in this letter, the problem has not been resolved yet.

The electron mass operator in a strong magnetic field with the one-loop contribution taken into account obtained by Jancovici [1] in the leading log approximation can be written as

$$M^{(1)} = m \left(1 + \frac{\alpha}{4\pi} \ln^2 \frac{eB}{m^2}\right). \tag{1}$$

The origin of the double logarithm asymptotics can be easily illustrated by using of the electron propagator in a strong magnetic field [2] for presenting the one-loop contribution in the form of the following integral

$$\Delta M^{(1)} = -\frac{i\alpha}{2\pi^3} m \int d^2k_\perp e^{-k_\perp^2/2eB} \int \frac{d^2k_\parallel}{[(k-p)_\parallel^2 - m^2](k_\parallel^2 - k_\perp^2)} \Big|_{p_\parallel^2 = m^2}, \tag{2}$$

where the notations are used: $k_\perp^2 = k_1^2 + k_2^2$, $k_\parallel^2 = k_0^2 - k_3^2$, (the magnetic field is directed along the 3d axis). The Wick rotation in the complex plane of k_0 , $k_0 = i\tilde{k}_0$ yields $k_\parallel^2 = -\tilde{k}_0^2 - k_3^2 \equiv -\tilde{k}_\parallel^2$. With the main contribution into the integral coming from the region $k_\perp^2 \gg \tilde{k}_\parallel^2 \gg m^2$, one obtains

$$\Delta M^{(1)} \simeq \frac{\alpha}{2\pi} m \int\limits_{m^2}^{\infty} \frac{dk_\perp^2}{k_\perp^2} e^{-k_\perp^2/2eB} \int\limits_{m^2}^{k_\perp^2} \frac{d\tilde{k}_\parallel^2}{\tilde{k}_\parallel^2}. \tag{3}$$

It is seen that the result contains the logarithm squared, and coinsides with the second term of eq. (1).

Later on, in the papers by Loskutov and Skobelev [3, 4] the attempts were performed to calculate the two-loop contribution and to summarize

[1] We use natural units in which $c = \hbar = 1$ and the pseudo-Euclidean metric with a $(+---)$ signature. Hereafter e is the elementary charge.

all the many-loop contributions in the same leading log approximation. A rather effective technique of "two-dimensional electrodynamics" in a strong magnetic field, developed by the authors was used and it was shown that the main contribution into the electron mass operator arises from the rainbow Feynman diagram depicted in Fig. 1. The result of summation over all orders of the perturbation theory was obtained in Ref. [4] in the exponential form:

$$M_{LS} = m \, \exp\left(\frac{\alpha}{4\pi} \, \ln^2 \frac{eB}{m^2}\right). \tag{4}$$

However, in the recent paper by Gusynin and Smilga [5] the summation over all orders of the perturbation theory was remade giving another result:

$$M_{GS} = m \Big/ \cos\left(\sqrt{\frac{\alpha}{2\pi}} \, \ln \frac{eB}{m^2}\right). \tag{5}$$

Obviously, the double logarithmic terms are also collected here because x^2 is the parameters of the $\cos x$ expansion. The first terms of expansion over α of both equations, (4) and (5), coinside and reproduce the one-loop result (1) of the paper [1]. The difference manifests itself at the two-loop level where the corresponding contributions from Eqs. (4) and (5) have the form

$$\Delta M^{(2)}_{LS} = m \, \frac{1}{32} \, \frac{\alpha^2}{\pi^2} \, \ln^4 \frac{eB}{m^2}, \tag{6}$$

$$\Delta M^{(2)}_{GS} = m \, \frac{5}{96} \, \frac{\alpha^2}{\pi^2} \, \ln^4 \frac{eB}{m^2}. \tag{7}$$

From the computation standpoint, in the frame of the approximation used by the authors, we confirm the result (7). It is intriguing to compare the Eqs. (6) and (7) with the earlier result for the two-loop contribution calculated in Ref. [3], see Eq. (5) of that paper. The numerical coefficient presented there, 5/48, was erroneous, to be replaced to 1/32, as it was pointed out in the next paper [4]. In fact, the true coefficient appeared to be 5/96.

However, both equations (1) and (7) had a restricted area of application because they were obtained without taking account of a crucial influence of the strong magnetic field on the virtual photon polarization operator. As was shown for the first time in Ref. [6], due to the photon polarization in the field, the double logarithm asymptotics is only valid at $B \lesssim B_e/\alpha$. For the larger field values, a photon acquires the effective mass $m_\gamma^2 = (2\alpha/\pi)eB$. Turning back to Eq. (2), we should replace the factor $k^2 = k_\parallel^2 - k_\perp^2$ in the

denominator arising from the massless photon propagator, to the expression $k^2-m_\gamma^2$. It changes the result essentially, because in one of the two logarithms the electron mass is replaced to the photon mass,

$$\ln(eB/m_\gamma^2) \sim \ln(1/\alpha),$$

so, the dependence of the electron mass operator on the field becomes not double but single logarithmic:

$$\Delta M^{(1)} = m\,\frac{\alpha}{2\pi}\,\left(\ln\frac{\pi}{\alpha}-\gamma_E\right)\,\ln\frac{eB}{m^2}, \tag{8}$$

where $\gamma_E = 0.577\ldots$ is the Euler constant.

However, the influence of the strong magnetic field on the photon polarization properties should be considered in more details. As was shown by Shabad, see, e.g. [7] where the earlier references can be found, the photon propagator in a magnetic field can be presented in the form

$$\mathcal{D}_{\mu\nu}(k) = -i\sum_{\lambda=1}^{3}\,\frac{1}{k^2-\mathcal{P}^{(\lambda)}}\,\frac{b_\mu^{(\lambda)}b_\nu^{(\lambda)}}{\left(b^{(\lambda)}\right)^2}, \tag{9}$$

where $b_\mu^{(\lambda)}$ are the eigenvectors and $\mathcal{P}^{(\lambda)}$ are the eigenvalues of the photon polarization operator,

$$b_\alpha^{(1)} = (q\varphi)_\alpha, \quad b_\alpha^{(2)} = (q\tilde{\varphi})_\alpha, \quad b_\alpha^{(3)} = q^2(q\varphi\varphi)_\alpha - q_\alpha\,(q\varphi\varphi q),$$

$\varphi_{\alpha\beta} = F_{\alpha\beta}\Big/\sqrt{F_{\mu\nu}^2/2}$ is the dimensionless tensor of the external magnetic field, $\tilde{\varphi}_{\alpha\beta} = \frac{1}{2}\varepsilon_{\alpha\beta\mu\nu}\varphi_{\mu\nu}$ is the dual tensor. The indices of four-vectors and tensors inside the parentheses are contracted consecutively, e.g. $(q\varphi)_\alpha = q_\beta\varphi_{\beta\alpha}$, etc.

It is well-known that only the photons, both real and virtual, with the $\lambda = 2$ polarization participate in all the electron - photon processes in a strong magnetic field. Taking into account that the electron mass being the physical value is gauge independent, it is convenient to calculate it in the gauge where the photon propagator of the mode 2 takes the form

$$\mathcal{D}_{\mu\nu}^{(2)}(k) = -i\,\mathcal{D}(k^2)\,\tilde{\varphi}_{\mu\rho}\tilde{\varphi}_{\rho\nu}, \quad \mathcal{D}(k^2) = \frac{1}{k^2-\mathcal{P}^{(2)}}. \tag{10}$$

To calculate the electron mass operator in the leading log approximation, it is enough to know the mode 2 photon polarization operator at the one-loop level [7]:

$$\mathcal{P}^{(2)} = -\frac{2\alpha}{\pi}\, eB\, H\left(\frac{k_\parallel^2}{4m^2}\right) + \frac{\alpha}{3\pi}\, k^2\, \ln\frac{eB}{m^2}. \tag{11}$$

Here, the first term arises from virtual electrons at the ground Landau level, while the second term is determined by the higher Landau levels. The function $H(k_\parallel^2/4m^2)$ for $k_\parallel^2 < 0$ has the form

$$H(z) = \frac{1}{2\sqrt{z(z+1)}} \ln\frac{\sqrt{z+1}+\sqrt{z}}{\sqrt{z+1}-\sqrt{z}} - 1, \quad z = \frac{|k_\parallel^2|}{4m^2}. \tag{12}$$

For the region of parameters we are interested in, $|k_\parallel^2| \gg 4m^2$, we have $H(z) \simeq -1$. Consequently, the first term in Eq. (11) acquires the meaning of the effective photon mass squared, $m_\gamma^2 = (2\alpha/\pi)eB$, induced by a magnetic field. Substituting (11) into Eq. (10), and multiplying by α, we obtain for the function $\alpha\,\mathcal{D}(k^2)$:

$$\alpha\,\mathcal{D}(k^2) = \frac{\alpha}{k^2 - k^2\,(\alpha/3\pi)\,\ln(eB/m^2) - m_\gamma^2(\alpha)}. \tag{13}$$

It is seen from Eq. (13) that the contribution of the higher Landau levels leads in fact to the electron charge renormalization in a strong magnetic field

$$\alpha \longrightarrow \alpha_R = \frac{\alpha}{1 - (\alpha/3\pi)\,\ln(eB/m^2)}. \tag{14}$$

Based on this, the Eq. (13) can be presented in the form

$$\alpha\,\mathcal{D}(k^2) = \frac{\alpha_R}{k^2 - m_\gamma^2(\alpha_R)}. \tag{15}$$

By this means the mode 2 photon manifests itself in a strong magnetic field as a massive vector boson interacting with an electron with the renormalized coupling constant α_R.

For the n-loop contribution into the electron mass operator in the leading log approximation, which is defined by the rainbow diagram, we have obtained

$$\Delta M^{(n)} = m\,\left[\frac{\alpha_R}{2\pi}\,\left(\ln\frac{\pi}{\alpha_R} - \gamma_E\right)\,\ln\frac{eB}{m^2}\right]^n. \tag{16}$$

Finally, for the electron mass operator in a strong magnetic field one obtains

$$M = m \Big/ \left[1 - \frac{\alpha_R}{2\pi} \left(\ln \frac{\pi}{\alpha_R} - \gamma_E\right) \ln \frac{eB}{m^2}\right]. \tag{17}$$

It should be noted that our expression (17) for the electron mass operator in the single logarithm field asymptotics is different from the analogous result obtained in Ref. [6], which had the form

$$M = m \ \exp\left(\frac{\alpha}{2\pi} \ln \frac{\pi}{\alpha} \ln \frac{eB}{m^2}\right). \tag{18}$$

To the first order in the parameter $\alpha \ln(eB/m^2)$, the results (17) and (18) almost coinside [2] and are reduced to our formula (8). However, already in the second order in $\alpha \ln(eB/m^2)$ the results are different because of two reasons. First, the arithmetic error was made in Ref. [6] in the calculation of the n-loop rainbow diagram which led to an incorrect summation of the perturbation expansion. Second, the authors [6] did not take into account the contribution of the higher Landau levels which is essential in the logarithm asymptotics. The fundamental difference of our result (17) from the result (18) of Ref. [6] is in the fact that the electron mass as the function of the field strength, $M(B)$, is the singular one, because the denominator of Eq. (17) can, in principle, tend to zero. It should be noted, however, that the singular behaviour of the function $M(B)$ could manifest itself only in a superstrong field $B \sim 10^{75}$ G.

Acknowledgements

We are grateful to V.A. Rubakov and to the participants of the seminar of the Theoretical Department of the Institute for Nuclear Research of RAS for fruitful discussion.

This work was supported in part by the Russian Foundation for Basic Research under the Grant N 01-02-17334 and by the Ministry of Education of Russian Federation under the Grant No. E00-11.0-5.

[2] The Euler constant γ_E is lost in Ref. [6].

References

[1] B. Jancovici, *Phys. Rev.* **187**, 2275 (1969).

[2] Yu. M. Loskutov, V. V. Skobelev, *Phys. Lett.* **A 56**, 151 (1976).

[3] Yu. M. Loskutov, V. V. Skobelev, *Teor. Mat. Fiz.* **38**, 195 (1979).

[4] Yu. M. Loskutov, V. V. Skobelev, *Teor. Mat. Fiz.* **48**, 44 (1981).

[5] V. P. Gusynin, A. V. Smilga, *Phys. Lett.* **B 450**, 267 (1999).

[6] Yu. M. Loskutov, V. V. Skobelev, *Vestn. Mosk. Univ., Fiz., Astron.* **24**, 95 (1983).

[7] A. E. Shabad, *Tr. Fiz. Inst. Akad. Nauk SSSR* **192**, 5 (1988).



Figure 1: The rainbow graph.